



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

0 253 190
A2

EUROPEAN PATENT APPLICATION

Application number: 87109318.3

Int. Cl. 4: C07K 5/02, C07K 1/00,
A61K 37/02, A61K 37/64

Date of filing: 29.06.87

Priority: 16.07.86 IT 2114586
16.07.86 IT 2114386

Date of publication of application:
20.01.88 Bulletin 88/03

Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

Applicant: ENIRICERCHE S.p.A.
Corso Venezia 16
I-20121 Milan(IT)

Applicant: SCLAVO S.p.A.
Via Fiorentina 1
I-53100 Siena(IT)

Inventor: Verdini, Antonio Silvio
Via S. Martino 12
I-Monterotondo(Roma)(IT)
Inventor: Bonelli, Fabio
Via Don Carlo Gnocchi 10
I-Roma(IT)
Inventor: Pessi, Antonello
Via Massaciuccoli 19
I-Roma(IT)
Inventor: Cardinalli, Franco
V. le Vega 18
I-Ostia Lido(Roma)(IT)
Inventor: Boraschi, Diana
Chiesa S. Leonardo a Catignano Pianella
I-Castelnuovo Berardenga(Siena)(IT)
Inventor: Censini, Stefano
Via Della Torre 7
I-Serre di Rapolano (Siena)(IT)
Inventor: Di Trapani, Romano
Via Corsica, 28
I-Monterotondo (Roma)(IT)


Representative: De Carli, Eida et al
ENIRICERCHE S.p.A. BRELD Via F. Maritano,
26
I-20097 San Donato Milanese(IT)

Partially retro-inverted tuftsin analogues, method for the preparation thereof and pharmaceutical compositions containing them.

New partially retro-inverted tuftsin analogues of general formula I

BEST AVAILABLE COPY

EP 0 253 190 A2



wherein

R represents the side-chain of the amino acids threonine, methionine or leucine

R¹ represents the side-chain of the amino acids lysine or arginine


R² is hydrogen or a metabolically labile acyl group,

all the asymmetric carbon atoms are either of the S- or R-configuration, or, alternatively, the first, third, and fourth asymmetric carbons, starting from the N-terminal residue, are of the S-configuration while the second one is of the R- or (R,S)-configuration, and the corresponding pharmacologically acceptable salts, esters and amides.

The new compounds which share the same pharmacological properties of tuftsin, are much more stable toward the enzymatic degradation than the parent molecule.

PARTIALLY RETRO-INVERTED TUFTSIN ANALOGUES, METHOD FOR THE PREPARATION THEREOF AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

The present invention refers to new partially retroinverted tuftsin analogues of general formula I



75 wherein

R represents the side-chain of the amino acids threonine ($R = -CH(OH)CH_3$), methionine ($R = -CH_2CH_2S-CH_3$) or leucine ($R = -CH_2CH(CH_3)_2$).

R¹ represents the side-chain of the amino acids lysine (R¹ = -(CH₂)₄-NH₂) or arginine (R¹ = -(CH₂)₃-NH-C(=NH)NH₂)

R₂ is hydrogen or a metabolically labile acyl radical.

and all the asymmetric carbon atoms are of the same S- or R-configuration, or the first, third, and fourth asymmetric carbons starting from the N-terminal residue are of the S-configuration while the second is of the R- or (R,S)-configuration.

and the corresponding pharmacologically acceptable salts, esters and amides,

For the purposes of the present invention, the term "metabolically labile acyl radical", as used above, designates any acyl radical which is rapidly cleaved off in the first steps of the metabolic pathway and which is relatively non-toxic and innocuous to mammals at dosages consistent with good biological activity.


A preferred group of compounds of the present invention comprises those compounds of formula I wherein R and R¹ are as defined above and R² is a hydrogen atom, and the corresponding pharmacologically acceptable salts, esters and amides.

A particularly preferred group of compounds of the invention comprises those compounds of formula I wherein R is as defined above, R¹ represents the side-chain of the amino acid lysine and R² is hydrogen, and the corresponding pharmacologically acceptable salts, esters, and amides.

A most preferred group of compounds of the invention comprises those compounds of formula I wherein R represents the side-chain of the amino acid threonine, R¹ represents the side-chain of the amino acid lysine, and R² is hydrogen, and the corresponding pharmacologically acceptable salts, esters and amides. Tuftsin is a natural tetrapeptide, characterized by the sequence Thr-Lys-Pro-Arg, which has been isolated in 1970 by Najjar and coworkers (see Najjar V.A. and Nishioka K., *Nature*, Vol.228, p.672 (1970)).

It is freed from a particular immunoglobulin, leukokinin, by the action of two enzymes, tuftsin-endocarboxypeptidase, a spleen enzyme which acts upon circulating leukokinin and cleaves the Arg-Glu bond at the carboxyterminal of tuftsin, and leukokininase, a membrane enzyme of neutrophils, monocytes and macrophages which cleaves the Lys-Thr bond thus setting free the aminoterminal of tuftsin. Its principal biological effect is to activate phagocytic cells, principally the macrophages, but it is fully active only when it is set free from the carrier leukokinin molecule (Najjar V.A. - *Advances in Enzymology* - 41 - p.129-78 (1974); Najjar V.A. - *J. Pediatr.* - 87 -p.1121-24 (1975)). Tuftsin binds to specific receptors on the outer membrane of phagocytic cells, after which it is internalized and becomes susceptible to the action of cytoplasmic enzymes. The most active enzyme is an aminopeptidase that cleaves off the threonine residue to yield the tripeptide Lys-Pro-Arg, which is an inhibitor of tuftsin activity (Spirer Z. et al. - *J.Clin.Invest.*, 55, p.198-200 (1975) and Fridkin M. et al. - *Biochim.Biophys.Acta*, 496, p.203-11 (1977)). Since 1970, and particularly in the last ten years, tuftsin has been thoroughly investigated.

in the presence of a suitably selected coupling agent, followed by reaction of the thus obtained compound of formula IV



wherein

R', R¹, P, and P' as well as the configuration of the asymmetrical carbon atoms are as defined above, with (1,1-bis-trifluoroacetoxy)iodobenzene (TIB) to selectively convert the terminal carbamyl group to a primary amino group, followed by optional acylation of the terminal amino function with a metabolically labile acyl group, and removal of the protecting groups. If a compound of formula I wherein R² is hydrogen is desired and the acylation step is therefore no longer needed, the sequential arrangement of the amide/amine conversion and deprotection steps may suitably be reverted. According to a preferred embodiment of the invention, in fact, when a compound of formula I is desired wherein R² is a hydrogen atom, the compound of formula IV, obtained by condensing fragments II and III, is deprotected first and then reacted with TIB to give the desired retro-inverso peptide of formula I.

The new intermediates of formula IV, wherein R', R¹, P, and P' are as defined above, as well as the corresponding derivatives wherein the protecting groups have been suitably removed, represent a further object of the present invention. In particular, said products, besides being useful as intermediates, proved to have an immunostimulating activity comparable to that of tuftsin itself.

As for a detailed description of the above sketched process, the first step may conveniently be carried out according to any of the peptide synthesis methods known in literature. Optimum results, in terms of yields and purity of the products, have been obtained by preparing first an activated ester of the carboxylic acid II, e.g. by the addition of a slight excess of N-hydroxy-benzotriazole (HOBT) to a solution of the acid of formula II, contacting then the activated ester with the coupling agent, typically dicyclohexylcarbodiimide (DCCI), and finally with the reaction partner of formula III. Conventional polar aprotic organic solvents capable of dissolving the reactants and which do not interfere with the reaction course, are suitably employed in this condensation reaction, which can be carried out conveniently at room temperature. Solvents of choice are the halogenated hydrocarbons, e.g. methylene chloride, chloroform, dichloroethane, etc., optionally mixed with more polar organic solvents such as dimethylformamide, dimethylsulfoxide, acetonitrile, and the like. When the condensation reaction, whose course can be easily monitored by tic, is complete, the intermediate product of formula IV can be recovered according to conventional recovery techniques. In particular, when DCCI is used as the coupling agent, these involve evaporating off the solvent, dissolving the residue in tetrahydrofuran, cooling the reaction mixture, removing the cyclohexylurea precipitate by filtration, carefully washing the filtrate with slightly basic and slightly acidic aqueous solutions and evaporating off the organic solvent.

The reaction of the thus obtained product with TIB is then carried out according to the method described in Italian patent application 25755 A/81, which involves reaction of the amide substrate with a slight excess of TIB in water/inert organic solvent mixtures such as for instance water/dimethylformamide, water/acetonitrile, and the like. The reaction is carried out while bubbling an inert gas through the reaction mixture and monitoring the reaction course by tic.

At the end of the reaction, the organic solvent is removed and the product is easily recovered by freeze-drying. Acylation of this product can then be carried out, using the active esters of the acid R₂COOH such as for instance the p-nitrophenyl or 2,4,5-trichlorophenyl esters. Deprotection is then carried out according to known methods. In general, when conventional protecting groups are employed, such as t-butyl or t-amyl groups for carboxyl protection and optionally for threonine hydroxyl protection, and tert-butoxycarbonyl, benzyloxycarbonyl and the like groups are employed for amino protection, these groups


are conveniently removed by acidolysis in mildly acidic conditions such as for instance with diluted hydrochloric acid in acetic acid, with trifluoroacetic acid or with trifluoroacetic acid/methylene chloride mixtures in the presence of varying percentages of anisole, thioanisole, or resorcinol used as scavengers to entrap the t-butyl, t-amyl or benzyl carbocations which form.

5 Once deprotection is over, the desired product of formula I is recovered and purified according to conventional chromatographic methods and techniques. Particularly suitable to this purpose is reversed-phase chromatography.

Homogeneity of the thus obtained products is checked by tlc and HPLC, while their purity is further tested by amino acid analysis and NMR spectroscopy.

10 As anticipated above, when a compound of formula I is desired wherein R² is hydrogen, the sequential arrangement of the amide/amine conversion and deprotection steps is preferably reverted. The general methods for carrying out each step are however the same.

The starting compounds of formulas II and III may be easily prepared either starting from commercially available compounds or from compounds prepared on purpose by methods known or analogous to methods known in peptide and organic synthesis chemistry. More particularly, the malonyl derivative of formula II is prepared by a series of condensation/deprotection steps carried out according to methods and techniques known in literature and widely employed in the synthesis of partially retro-inverted peptides, starting from the malonyl derivative V




wherein R^{1'} is as defined above.

25 In its turn the malonyl derivative V is easily prepared from the corresponding di-ester by partial hydrolysis under controlled alkaline conditions. The diester is obtained through reaction of the alkaline salt of a malonic acid di-lower alkyl ester with a suitably selected halide of formula R^{1'}-X wherein R^{1'} is as defined above and X is preferably chloro. More particularly, the malonic acid di-lower alkyl ester is added to an alcohol solution of the alkali metal, and the reaction partner R^{1'}-X is then added thereto. When the reaction is complete, the desired product is recovered by washing with water and evaporating off the organic solvent.

Partial hydrolysis of the thus obtained intermediate, typically in ethanol/KOH, affords the mono-ester of formula V.

35 As the compounds of the present invention have at least 4 asymmetric carbon atoms in their molecule (those marked by an asterisk in the formula below), plural isomeric forms may exist



I

50 The above summarized process, however, is a stereo-selective process i.e. the configuration of each starting fragment, excluding the 2-substituted malonyl residue which is employed as a racemate, is retained in each step.

55

According to a preferred embodiment of the present invention the absolute configuration of the terminal amino acids Pro-Arg is the L-and the compounds of formula I wherein the last two asymmetric carbon atoms are of the L-configuration can be easily obtained using the suitable L-proline and L-arginine derivatives in the preparation of the starting compound of formula II. As the use of the 2-substituted malonyl residue in the above defined process yields a mixture of isomers of formula I, this can be resolved, if desired, into the single isomers, generally by reversed-phase HPLC.

The compounds of the present invention of formula I may form basic and acid salts with a number of inorganic and organic bases and acids. Said basic salts include for instance the alkali metal salts, such as sodium or potassium, or earth-alkaline salts such as calcium or magnesium, while the acid salts include the acid addition salts with inorganic acids such as HCl, HBr, H_2SO_4 , H_3PO_4 , sulphonic organic acids and carboxylic organic acids such as acetic acid, oxalic acid, pivalic acid and the like acids. The physiologically acceptable salts are obviously preferred as they can be employed in therapy as described later on. These and other salts, which are not necessarily physiologically acceptable, may however be employed for isolating or purifying the compounds of formula I. Said salts are easily prepared by reacting the compound of formula I as the free acid or base with one or more equivalents of the suitably selected base or acid in a solvent or reaction medium in which the salt is insoluble or in a solvent which can be easily removed. Following a particularly advantageous technique, said salt formation is carried out in water and the thus obtained salt is recovered by freeze-drying.

The corresponding esters and amides of the compounds of formula I are also enclosed within the scope of the present invention. Said derivatives include the alkyl, di-(alkyl)aminoalkyl, acylaminoalkyl, acyloxyalkyl, benzyl and substituted benzyl esters and the benzyl, phenethyl, and mono- and di-N-alkyl amides. A preferred group of said derivatives includes the esters of the compounds of formula I wherein the ester group is easily hydrolysed in acidic conditions, such as for instance the t-butyl or t-amyl ester, because these compounds can be used as intermediates.

The compounds of formula I, including the pharmaceutically acceptable esters, amides and salts, are active as immunostimulating agents. This activity has been demonstrated *in vitro* by means of the macrophage (M ϕ) phagocytic stimulation assay. This assay, carried out in comparison with tuftsin, has been performed according to a modification of the method described by Bar-Shavit et al. in J.Cell.Physiol., 100, p.55 (1979). More particularly, peritoneal M ϕ from normal C3H/HeN mice are harvested by washing of the peritoneal cavity and purified by adherence as described in J.Immunol., 132, p.1987 (1984). Monolayers of 1×10^6 adherent peritoneal cells (95-98% of the monocyte-M ϕ series as judged by morphology and latex phagocytosis) are cultured for 20 h at 37 °C in RPMI-1640 medium supplemented with 50 μ g/ml gentamycin, 25 mM HEPES buffer, 2 mM L-glutamine and 10% heat-inactivated fetal bovine serum (FBS). At the end of the incubation, M ϕ are washed and then exposed for 15 minutes at 37°C to 1.3 ml/plate of culture medium either alone (blank) or containing tuftsin or a compound of the invention at different concentrations. A 1.5×10^5 cells/ml suspension of killed yeast cells (Zymosan)(0.2 ml) is then added to the plates. M ϕ are incubated for 60 minutes at 37°C, then washed, fixed (2% glutaraldehyde in PBS, 30 minutes, 4°C) and stained. The percentage of phagocytosing M ϕ and the number of ingested Zymosan cells is measured with an optical microscope equipped with a 100x immersion objective. In a representative experiment aimed at comparing the activity of tuftsin with that of the corresponding retro-inverso analogue of example 1, it has been observed that all M ϕ were capable of phagocytosing zymosan in the above described conditions. However, in the presence of tuftsin or of the compound of example 1, the number of ingested yeast cells significantly increased and was linearly dependent on peptide concentration. In particular it has been observed that the compound of example 1 at a concentration of 3×10^{-7} M resulted in 2.2×10^3 ingested zymosan cells per 100 macrophages (170 % of the control), while at the concentration of 10^{-6} M gave 2.44×10^3 ingested zymosan cells per 100 phagocytosing macrophages (188% of the control).

As the poor therapeutical applicability of tuftsin is mainly due to its low stability and to the consequent rapid loss of biological activity, the phagocytosis assay has been repeated comparing the activity of tuftsin and its retro-inverso analogue of example 1, stored for 14 days in solution at 4°C. In these conditions tuftsin, as expected, was devoid of activity, while, on the contrary, the phagocytosis stimulation induced by the compound of example 1, was still significant even if less than above. At the concentration of 10^{-6} M in fact the compound of Example 1 gave a number of ingested zymosan cells per 100 phagocytosing macrophages of 1.8×10^3 (with an immuno-potentiating effect corresponding therefore to 136 % of the control).

The compounds of the present invention are therefore useful in all those cases where it is necessary or advisable to stimulate the defense mechanisms against infectious agents or tumor cells. For said pathological conditions, a treatment schedule providing for a single dose, repeated every 4-8 days, can conveniently be employed where the dosage, which obviously depends on the pathological condition and its severeness,

the administration route, the particular compound of formula I administered and any concomitant therapeutic treatment, will be generally comprised between 0.001 and 5 mg of active principle per Kg of body weight and preferably between 0.01 and 2 mg/Kg. The compounds of formula I can be administered parenterally, and in particular intravenously and intraperitoneally, or orally. They can therefore be formulated in suitable solid or liquid dosage forms, such as for instance tablets, capsules or elixirs for the oral administration and sterile solutions or suspensions for parenteral administration.

Said dosage forms, which will preferably contain from 0.01 to 300 mg of a compound of formula I as the active ingredient per unit dosage form, together with the conventional vehicles, excipients, preservatives, and the like agents, are prepared according to methods well known to the industrial pharmacy expert.

The compounds of the present invention may also be formulated in admixture with other therapeutically effective compounds. For instance, in the treatment of bacterial infections in patients with impaired immune functions, it may be advisable to use a formulation containing, besides the antibiotic agent effective against the microorganism causing the infection, a suitable dose of an immunostimulating compound of formula I.

The following example which is only aimed at better describing the preparation of a representative compound of the invention, should in no way be considered as a limitation to the scopes thereof.

For the sake of clarity, the following abbreviations have been used throughout the text.

Z = benzyloxycarbonyl; Bu^t = tert-butyl; Boc = tert-butoxycarbonyl; AcOEt = ethyl acetate; Mtr = (2,3,6-trimethyl-4-methoxyphenyl)sulphonyl; EtO = ethoxy; Bz = benzyl; NMM = N-methylmorpholine; HOBT = N-hydroxybenzotriazole; DMF = dimethylformamide; DCCI = dicyclohexylcarbodiimide; EtOH = ethyl alcohol; THF = tetrahydrofuran; DCU = dicyclohexylurea; TFA = trifluoro-acetic acid; Et₂O = ethyl ether; TIB = (1,1-bis-trifluoroacetoxy)iodobenzene;

Example 1

{(2RS)-2-[N-(1-amino-2-hydroxy-propyl)carbonyl]-6-amino}-hexanoyl-L-prolyl-L-arginine

gThr-(R,S)mLys-L-Pro-L-Arg-OH

a) N-benzyloxycarbonyl-O-tert-butyl-D-threonineamide (Z-D-Thr(Bu^t)-NH₂)

Concentrated H₂SO₄ (0.1 ml) and isobutylene (35 ml, 390 mmol) are added to a suspension of N-benzyloxycarbonyl-D-threonineamide (4.2 g, 16.6 mmol) in methylene chloride (35 ml) kept in a pressure resistant vessel cooled with dry ice/acetone. The reaction vessel is sealed and the temperature is allowed to raise up to the room value. After 4 days, excess isobutylene is evaporated off and the organic solution is washed with aqueous 5% Na₂CO₃ (3 × 30 ml), 5% citric acid (20 ml) and then with water up to pH 6. The organic phase is dried over MgSO₄ and concentrated to dryness yielding the compound of the title as a clear oil (4.62 g, 90%).

NMR and mass spectroscopy confirm the assigned structure. HPLC and tlc confirm that the product is unitary.

b) O-tert-butyl-D-threonineamide (H-D-Thr(Bu^t)-NH₂)

10% Pd/C (1.5 g) is added to a solution of the compound obtained in step a) in MeOH (200 ml), and hydrogen is bubbled through the reaction mixture for 1.5 h, checking the disappearance of the starting compound by tlc. When the reaction is over, excess hydrogen is removed by passing a nitrogen stream through the mixture, the reaction mixture is filtered over celite and the filtrate is concentrated to dryness thus yielding O-tert-butyl-D-threonineamide (2.55 g, 99%). Both mass and NMR spectra confirm the assigned structure.

c) mono-ethyl ester of (2RS)-2-[(4-t-butoxy-carbonyl-amino)butyl]malonic acid

OEI-(R,S)mLys(Boc)-OH

A solution of 4-chloro-butylamine hydrochloride (14.38 g, 0.1 mol) in dioxane/water (100 ml, 2/1, v/v) is slowly added to a vigorously stirred mixture of di-(tert-butyl)carbonate (24 g, 0.11 mol), 1N Na₂CO₃ (100 ml) and dioxane/water (200 ml, 2/1, v/v) cooled to 0°C. When the addition is over, the reaction mixture is stirred at room temperature for an additional hour, dioxane is removed under vacuum and the aqueous phase is extracted a few times with ethyl acetate. From the organic solution N-[(tert-butoxy)carbonyl]-4-chloro-butylamine, as an oily product (21 g), is recovered. Sodium metal (0.28 g, 0.012 mol) is dissolved in absolute ethyl alcohol (9 ml) under nitrogen atmosphere. The mixture is heated to 60°C and malonic acid di-ethyl ester (3.8 g, 0.024 mol) is slowly dripped in.

N-[(tert-butoxy)carbonyl]-4-chloro-butyl-amine (2.5 g, 0.012 mol) is then gradually added to the resulting mixture at room temperature. The reaction mixture is stirred at room temperature for 2 hours and at the reflux temperature for 6 hours, then it is poured into ethyl acetate/water (100 ml, 1/1, v/v). The organic phase is recovered, washed several times with water and dried over MgSO₄. The organic solvent is removed under vacuum (0.5 mBar) at 100°C, yielding a raw oily product which is purified by reversed-phase HPLC using an RP-18 resin and eluting with an aqueous phase modified With CH₃CN (45 % by volume).

(2 R,S)-2-[(4-tert-butoxy-carbonyl-amino)butyl]-malonic acid diethyl ester (1.31 g) is thus obtained as a pure product.

An ethanol solution of KOH (4.31 ml, 0.417M) is very slowly added to a solution of (2R,S)-2-[(4-tert-butoxycarbonylamino)butyl]malonic acid diethyl ester (1.2 g, 3.6 mmol) in absolute ethyl alcohol (7 ml). The reaction mixture is allowed to stand overnight, then is diluted with water and ethanol is evaporated off. The pH is brought to 8 by the addition of a 5% NaHCO₃ solution and the reaction mixture is extracted with AcOEt (4 × 50 ml) in order to recover the unreacted starting compound. The aqueous solution is brought to pH 3 by the addition of citric acid and is extracted with AcOEt (6 × 50 ml). The organic extracts are combined, washed with water up to pH 6 and concentrated to dryness yielding the compound of the title as a colorless oil (1.6 g, 88%). The mass and NMR spectra are in agreement with the assigned structure.

d) N^α-benzyloxycarbonyl-N^ε-(2,3,6-trimethyl-4-methoxy-phenyl)sulphonyl-L-arginine tert-butyl ester
(Z-Arg(N^ε-Mtr)-OBu^t)

POCl₃ (7.06 ml, 77 mmol) is dripped into a solution of N^α-benzyloxycarbonyl-N^ε-(2,3,6-trimethyl-4-methoxyphenyl)sulphonyl-L-arginine (3.6 g, 7 mmol) and pyridine (6.94 ml, 86 mmol) in tert-butyl alcohol (100 ml) kept under nitrogen atmosphere and cooled to - 5 °C. The reaction mixture is allowed to stand at room temperature overnight and then it is diluted with a mixture of ethyl acetate and water 1/1. The organic phase is separated and washed first with a 5% NaHCO₃ aqueous solution (16 × 100 ml) and then with water up to neutral pH. The organic phase is dried over MgSO₄ and the solvent is evaporated off giving the compound N^α-benzyloxycarbonyl-N^ε-(2,3,6-trimethyl-4-methoxyphenyl)sulphonyl-L-arginine tert-butyl ester (1 g, 25%).

e) N^ε-(2,3,6-trimethyl-4-methoxy-phenyl)sulphonyl-L-arginine tert-butyl ester (H-Arg(N^ε-Mtr)-OBu^t)

By following substantially the same procedure as in step b) but starting from the compound obtained in step d), N^ε-(2,3,6-trimethyl-4-methoxyphenyl)-sulphonyl-L-arginine tert-butyl ester (612 mg, 81%) is obtained as a white foam.

[α]_D²⁰ -0.84° (c = 1.18% CH₂Cl₂)

The NMR spectrum confirms the assigned structure.

f) [(2RS)-2-ethoxy-carbonyl-6-tert-butoxy-carbonyl-amino]hexanoyl-L-proline benzyl ester

(EtO-(R,S)mLys(N^ε-Boc)-L-Pro-OBz)

The compound obtained in step c) (0.6 g, 1.98 mmol) is dissolved in CH₂Cl₂ (40 ml) and a solution of HOBT (0.325 g, 2.4 mmol) in DMF (2 ml) and CH₂Cl₂ (3 ml) is added thereto. The reaction mixture is cooled to 0°C and a solution of DCCl (0.495 g, 2.4 mmol) in CH₂Cl₂ (5 ml) is added thereto at this temperature. The reaction mixture is stirred at 0°C for 30 minutes and then at room temperature for additional 20 minutes. The activated ester thus prepared is filtered into a reaction flask containing a solution of proline benzyl ester hydrochloride (0.532 g, 2.2 mmol) in CH₂Cl₂ (60 ml), to which NMM (0.242 ml, 2.2 mmol) necessary to remove the hydrochloride has just been added. The mixture is stirred at room temperature overnight, the solvent is then evaporated off and the oily residue is taken up in a small amount of AcOEt and kept at -25°C for one hour. The precipitate is removed by filtration and the filtrate is diluted with an additional amount of AcOEt, washed with 5% aqueous NaHCO₃ (5 × 100 ml), with 5% aqueous citric acid (3 × 50 ml) and finally with water up to pH 6-6.5. The organic phase is dried over MgSO₄ and then concentrated to dryness under vacuum yielding [(2 R,S)-2-ethoxycarbonyl-6-tert-butoxycarbonyl-amino]hexanoyl-L-proline benzyl ester as a yellow oil (0.88 g, 90%). Mass and NMR spectra confirm the proposed structure.

g) [(2 R,S)-2-ethoxycarbonyl-6-(tert-butoxycarbonyl-amino)]-hexanoyl-L-proline20 (OEt-(R,S)mLys(N^ε-Boc)-L-Pro-OH)

This compound is prepared through hydrogenolysis of the compound obtained in the foregoing step, following substantially the same procedure as in step b).

The thus obtained product (91%) has $[\alpha]_D^{20} =$
 -38.18° (c = 1.32 % in CH₂Cl₂)

h) [(2R,S)-2-ethoxycarbonyl-6-(tert-butoxycarbonylamino)]hexanoyl-L-prolyl-(N^G-(2,3,6-trimethyl-4-methoxyphenyl)sulphonyl)-L-arginine tert-butyl ester(OEt-(R,S)mLys(N^ε-Boc)-L-Pro-L-Arg(N^G-Mtr)-OBu^t)

A solution of HOBT (0.261 g, 1.93 mmol) in CH₂Cl₂ (3 ml) and DMF (0.5 ml) is added to a solution of the compound prepared in step g) (0.66 g, 1.64 mmol) in CH₂Cl₂ (50 ml), the obtained mixture is cooled to 0 °C and DCCl (0.398 g, 1.93 mmol) is added thereto. The reaction mixture is then stirred at 0 °C for 30 minutes and at room temperature for further 20 minutes. The thus obtained activated ester is filtered into a reaction flask containing a solution of the compound of step e) (0.590 g, 1.3 mmol) in CH₂Cl₂ (60 ml). When the reaction, whose course can be easily monitored by tlc, is over, the solvent is evaporated off, the residue is taken up in a small amount of AcOEt and cooled to - 25°C for one hour. The precipitate is removed by filtration, the filtrate is diluted with additional AcOEt (100 ml) and washed with a 5% NaHCO₃ aqueous solution (4 × 100 ml), with a saturated NaCl solution and finally with water up to neutral pH. The organic solution is dried over MgSO₄ and the solvent is evaporated under vacuum yielding the desired compound (0.857 g, 80%) as a yellow oil.

i) [(2R,S)-2-carboxy-6-(tert-butoxycarbonylamino)] hexanoyl-L-prolyl-[N^G-(2,3,6-trimethyl-4-methoxyphenyl)-sulphonyl]-L-arginine tert-butyl ester50 (HO-(R,S)mLys(N^ε-Boc)-L-Pro-L-Arg(N^G-Mtr)-OBu^t)

1M KOH in absolute ethanol (2 ml, 2mmol) is added to a stirred solution of the compound obtained in the foregoing step (0.857 g, 1.04 mmol) in absolute ethanol (10 ml) cooled to 0°C, and the reaction mixture is stirred overnight. The mixture is then diluted with water, ethanol is evaporated off and the pH is brought to 3 by the addition of citric acid. The acidic mixture is extracted with AcOEt (4 × 60 ml), the organic

extracts are combined, washed with a saturated NaCl aqueous solution and then with water up to neutral pH, the organic phase is dried over MgSO_4 and the organic solvent is evaporated off yielding [(2 R,S)-2-carboxy-6-(tert-butoxy-carbonyl-amino)hexanoyl-L-prolyl = [N^G-(2,3,6-tri-methyl-4-methoxy-phenyl)-sulphonyl]-L-arginine tert-butyl ester (1.03 g).

5

j) [(2 R,S)-2-[N-(1-carbamyl-2-tert-butoxy-propyl)-carbamyl]-6-tert-butoxy-carbonyl-amino]hexanoyl-L-prolyl-(N^G-(2,3,6-trimethyl-4-methoxy-phenyl)-sulphonyl)-L-arginine tert-butyl ester

10 (H₂N-D-Thr(Bu^t)-(R,S)mLys(N^G-Boc)-L-Pro-L-Arg(N^G-Mtr)-OBu^t)

A solution of HOBT (0.194 g, 1.43 mmol) in DMF (1 ml) and CH_2Cl_2 (2 ml) is added to a solution of the compound obtained in the foregoing step (1.03 g, 1.3 mmol) in CH_2Cl_2 (30 ml) cleared by filtration. The reaction mixture is then cooled to 0°C and DCCl (0.268 g, 1.3 mmol) is added thereto. The mixture is stirred at 0 °C for 30 minutes and at room temperature for further 20 minutes. The mixture is filtered into a reaction vessel containing a solution of the compound obtained in step b) (0.333 g, 1.9 mmol) in CH_2Cl_2 (30 ml) and stirred overnight. The solvent is evaporated off to dryness, the residue is taken up in THF (10 ml) and the solution is cooled to -17°C for 2 hours. The DCU which precipitates is removed by filtration and the solution is brought to dryness yielding a clear, light yellow, oil which is dissolved in AcOEt (6 ml) and washed with 5% aqueous NaHCO_3 (4 × 100 ml), 5% aqueous citric acid and then with water up to neutral pH. The organic solution is dried over MgSO_4 and the solvent is evaporated off yielding a white foam (1.25 g). This product is purified by flash chromatography using a silica column and eluting with 60% n-hexane/acetone. The thus obtained product is freeze-dried and employed as such in the following step.

25

k) [(2R,S)-2-[N-(1-carbamyl-2-hydroxypropyl)carbamyl]-6-amino]hexanoyl-L-prolyl-L-arginine

(H₂N-D-Thr-(R,S)mLys-L-Pro-L-Arg-OH)

The compound obtained in the foregoing step is dissolved in TFA (20 ml) containing 6% thioanisole and the thus obtained solution is stirred at room temperature for 4 hours. TFA and thioanisole are then allowed to evaporate off by passing a nitrogen stream throughout the mixture, the residue is taken up in CH_3CN and concentrated to dryness under vacuum. The residue is taken up again in water containing a few drops of CH_3CN . It is washed with Et_2O (2 × 30 ml), Et_2O is evaporated off from the aqueous phase which is then freeze-dried.

The thus obtained raw product is purified by column chromatography eluting with $\text{CH}_3\text{COONH}_4$ $2 \times 10^{-2} \text{M}/\text{CH}_3\text{CN}$ 0.5%.

40 1) [(2R,S)-2-[N-(1-amino-2-hydroxypropyl)carbamyl]-6-amino]hexanoyl-L-prolyl-L-arginine

gThr-(R,S)mLys-L-Pro-L-Arg-OH

A solution of the compound obtained in the foregoing step (0.02 mmol) in DMF is added to a solution of TIB (0.024 mmol) in DMF/ H_2O 4/1 (10 ml) and the obtained reaction mixture is kept at 20°C under a nitrogen stream for 16 hours.

DMF is then removed and the aqueous phase is freeze-dried. The thus obtained product is purified by reversed-phase chromatography on a Lichroprep RP-18 column (40 g) eluting first with $\text{H}_2\text{O}/\text{TFA}$ 1%/CH₃CN 3% (540 ml) and then with $\text{H}_2\text{O}/\text{TFA}$ 1%/CH₃CN 75% (360 ml) and checking the eluted fractions by HPLC. Fractions containing only the desired product are combined. The product isolated therefrom, which as shown by tic and HPLC analyses is a unitary product, has the following amino acid composition (upon acid hydrolysis at 118°C for 18 hours) :


Thr	Pro	Arg
/	1.01	1.00

By following substantially the same procedure described in the foregoing example, or the general methods taught in the descriptive portion, and starting from the corresponding compounds of formulas II and III, the following compounds are obtained :

- 5 gThr-(R,S)mArg-Pro-Arg-OH
 gThr-(S)mArg-Pro-Arg-OH
 gThr-(S)mArg-Pro-Arg-OMe
 gThr-(S)mLys-Pro-Arg-ONa
 gMet-(R,S)mLys-Pro-Arg-OH
 10 gMet-(S)mLys-Pro-Arg-OH
 gLeu-(S)mLys-Pro-Arg-OH
 For-gLeu-(S)mLys-Pro-Arg-ONa
 For-gThr-(S)mLys-Pro-Arg-OH

Claims

1) A compound of general formula I



wherein

R represents the side-chain of the amino acids threonine, methionine or leucine,

R¹ represents the side-chain of the amino acids lysine or arginine, and

R² is a hydrogen atom or a metabolically labile acyl group.

and the corresponding pharmacologically acceptable salts, esters and amides.


2) A compound as in claim 1 wherein R² is hydrogen.

3) A compound as in claim 2 wherein R¹ represents the side-chain of the amino acid lysine.

4) A compound as in claim 3 wherein R represents the side-chain of the amino acid threonine.

5) A compound as in claim 4 which is the {2-[N-(1-amino-2-hydroxypropyl)carbonyl]-6-amino} hexanoyl-prolyl-arginine.

6) A process for preparing a compound of claim 1 which consists of condensing a malonic acid derivative of formula II



wherein


R^{1'} is the side-chain of the amino acids lysine or arginine wherein the amino or guanidino group is suitably protected,

P is an easily removable carboxyl protecting group
P' is an easily removable guanidino protecting group
with an amide of formula III



wherein


10 R represents the side-chain of the amino acids threonine, methionine or leucine wherein the functional groups, if any, are suitably protected,
in the presence of a suitable coupling agent, reacting the thus obtained compound of formula IV



25 wherein

30 R', R¹, P, and P' are as defined above, with (1,1-bis-trifluoroacetoxy)iodobenzene to convert the terminal amido group in primary amino, acylating, if desired, the terminal amino function with a metabolically-labile acyl group, removing the protecting groups and, if desired, converting the thus obtained compound of formula I into the corresponding pharmacologically acceptable salts, esters and amides.


7) A process for preparing a compound of claim 2 which consists in condensing a malonic acid derivative of formula II



45 wherein

R¹ is the side-chain of the amino acids lysine or arginine wherein the amino or guanidino group is suitably protected.

50 P is an easily removable carboxyl protecting group
P' is an easily removable guanidino protecting group
with an amide of formula III



wherein


R' represents the side-chain of the amino acids threonine, methionine and leucine wherein the functional

groups, if any, are suitably protected.

in the presence of a suitable coupling agent, removing the protecting groups, reacting the thus obtained compound with (1,1-bis-trifluoroacetoxy)-iodobenzene and, if desired, converting the thus obtained compound of formula I wherein R² is hydrogen into the corresponding pharmacologically acceptable salts, esters and amides.

8) A pharmaceutical composition comprising an immunostimulating effective amount of the peptides of claims 1, 2, 3, 4 or 5 alone or in combination With a pharmaceutically acceptable carrier.

9) A compound of general formula IV



wherein

R', R¹, P, and P' are as defined in claim 6 and the corresponding derivatives wherein the protecting groups have been removed.

10) A compound of claim 9 which is {2-[N-(1-carbamyl-2-hydroxy-propyl)carbamyl]-6-amino}hexanoyl-L-prolyl-L-arginine.

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

**0 253 190
A3**

12

EUROPEAN PATENT APPLICATION

21 Application number: 87109318.3

51 Int. Cl.⁴: C07K 5/02 , C07K 1/00 ,
A61K 37/02

22 Date of filing: 29.06.87

30 Priority: 16.07.86 IT 2114586
16.07.86 IT 2114386

43 Date of publication of application:
20.01.88 Bulletin 88/03

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

89 Date of deferred publication of the search report:
04.10.89 Bulletin 89/40

71 Applicant: ENIRICERCHE S.p.A.
Corso Venezia 16
I-20121 Milan(IT)

Applicant: SCLAVO S.p.A.
Via Fiorentina 1
I-53100 Siena(IT)


72 Inventor: Verdini, Antonio Silvio
Via S. Martino 12
I-Monterotondo(Roma)(IT)
Inventor: Bonelli, Fabio
Via Don Carlo Gnocchi 10
I-Roma(IT)
Inventor: Pessi, Antonello
Via Massaciuccoli 19
I-Roma(IT)
Inventor: Cardinali, Franco
V. le Vega 18
I-Ostia Lido(Roma)(IT)
Inventor: Boraschi, Diana
Chiesa S. Leonardo a Catignano Pianella
I-Castelnuovo Berardenga(Siena)(IT)
Inventor: Censini, Stefano
Via Della Torre 7
I-Serre di Rapolano (Siena)(IT)
Inventor: Di Trapani, Romano
Via Corsica, 28
I-Monterotondo (Roma)(IT)

73 Representative: Cioni, Carlo et al
c/o ENIRICERCHE S.p.A. BRELID Via F.
Maritano 26
I-20097 San Donato Milanese(IT)

EP 0 253 190 A3

54 Partially retro-inverted tuftsin analogues, method for the preparation thereof and pharmaceutical compositions containing them.

57 New partially retro-inverted tuftsin analogues of general formula I



wherein

R represents the side-chain of the amino acids threonine, methionine or leucine

R' represents the side-chain of the amino acids lysine or arginine

R² is hydrogen or a metabolically labile acyl group.

all the asymmetric carbon atoms are either of the S- or R-configuration, or, alternatively, the first, third, and fourth asymmetric carbons, starting from the N-terminal residue, are of the S-configuration while the second one is of the R- or (R,S)-configuration, and the corresponding pharmacologically acceptable salts, esters and amides.

The new compounds which share the same pharmacological properties of tuftsin, are much more stable toward the enzymatic degradation than the parent molecule.



EP 87 10 9318

DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A, D	CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 26, 1978, pages 1006-1007, Tokyo, JP; K. HISATSUNE et al.: "Structure-activity relationship of tuftsin, a phagocytosis-stimulating peptide, and its analogs" -----		C 07 K 5/02 C 07 K 1/00 A 61 K 37/02
A	EP-A-0 097 994 (ANIC S.p.A.) -----		
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 K A 61 K
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26-06-1989	Examiner DEFFNER C-A.E.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.